

Cereal Rust Report

2016

VOLUME 14 ISSUE 6

31st August



Wheat leaf rust and wheat stripe rust situation, late Winter 2016

DR WILLIAM CUDDY

Co-located at the NSW Department of Primary Industries, Elizabeth Macarthur Agricultural Institute, Menangle and The University of Sydney, Plant Breeding Institute, Cobbitty
Email: will.cuddy@dpi.nsw.gov.au Phone: 02-9351 8871

PROFESSOR ROBERT PARK

The University of Sydney, Plant Breeding Institute, Cobbitty
Email: robert.park@sydney.edu.au Phone: 02-9351 8806

This report provides a brief update on the current distribution of wheat leaf rust and wheat stripe confirmed from samples received at the Australian Cereal Rust Survey at the University of Sydney's Plant Breeding Institute. Details on the current distributions are provided as are pathotype details where known. Wheat growers are advised to inspect their crops for wheat leaf rust and wheat stripe rust and to submit samples of any rusts observed to the Australian Cereal Rust Survey. Submission details are provided on the back page of this report.

Wheat leaf rust

Samples of wheat leaf rust have been received from Victoria, South Australia and Queensland. Victorian samples off the wheat varieties SQP Revenue, Manning and Sunlamb were received from Bairnsdale in mid-June. The pathotypes identified from these samples included: 104-1,3,5,7,9,10,12+Lr37, 104-1,3,4,6,7,8,10,12+Lr37 and 76-1,3,5,7,9,10,12+Lr37. A sample off SQP Revenue was received from Lismore in western Victoria in late June and was determined to be pathotype 104-1,3,4,6,7,8,10,12+Lr37.

A sample of wheat leaf rust off Mace was received from Port Neill in South Australia in late June and a further sample off wheat from Paskeville was received in late August. Pathotype identifications are continuing for these samples.

Samples of wheat leaf rust off Sunzell were received from Millmerran and Mirabooka in Queensland in mid- and late-August respectively. Pathotype identifications for these two samples are underway. A sample of wheat leaf rust was also received from natural

infection at Cleveland and was determined to be pathotype 104-1,3,4,6,7,8,10,12+Lr37.

Samples of wheat leaf rust off Mace from Western Australia were received in mid-July from Coomalbidgup and late August from Grass Patch. Pathotype identifications are underway.

Pathotype 104-1,3,4,6,7,8,10,12+Lr37 was first detected in South Australia in 2014 and was the most recent exotic incursion of a wheat leaf rust pathogen into Australia. It has since spread across the eastern and western grainbelts and was a commonly sampled pathotype in the eastern grainbelt in 2015.

Wheat stripe rust

Samples of wheat stripe rust have been received from southern NSW and Queensland. Samples from Sandigo, Wagga Wagga and Colingullie in NSW were received from mid- to late-August. A sample off Suntop was received from Bungunya in Queensland in late August. Pathotype identifications are underway.

GENERAL ENQUIRIES

Mr Keshab Kandel
Rust Surveillance Technician
Plant Breeding Institute
Private Bag 4011,
Narellan NSW 2567

T 02-9351 8849
F 02-9351 8875

RUSTED PLANT SAMPLES

can be mailed in paper envelopes;
do not use plastic wrapping or plastic
lined packages. If possible, include the
latitude and longitude of the sample
location.

Direct samples to:

University of Sydney
Australian Rust Survey
Reply Paid 88076
Narellan NSW 2567

The Australian Cereal Rust Control Program is supported by growers through the Grains Research & Development Corporation.

GRDC **Grains Research & Development Corporation**
Your GRDC working with you



Department of Primary Industries