

2 Unit Bridging Course – Day 8


Composite Functions

Collin Zheng



The Composition of Two Functions

Consider two functions $f(x) = x + 1$ and $g(u) = u^2$.




A new function $(g \circ f)(x)$ can be formed from f and g by setting the output of f to be the input of g , giving $g(f(x)) = (x + 1)^2$.




$g \circ f$ is read as “ g composed with f ”, and the function $(g \circ f)(x) = g(f(x))$ is called a **composite function**.

The Composition of Two Functions

Consider two functions $f(x) = x + 1$ and $g(u) = u^2$.




A new function $(g \circ f)(x)$ can be formed from f and g by setting the output of f to be the input of g , giving $g(f(x)) = (x + 1)^2$.




$g \circ f$ is read as “ g composed with f ”, and the function $(g \circ f)(x) = g(f(x))$ is called a **composite function**.

The Composition of Two Functions

Consider two functions $f(x) = x + 1$ and $g(u) = u^2$.



A new function $(g \circ f)(x)$ can be formed from f and g by setting the output of f to be the input of g , giving $g(f(x)) = (x + 1)^2$.



$g \circ f$ is read as “ g composed with f ”, and the function $(g \circ f)(x) = g(f(x))$ is called a **composite function**.

Identifying Composite Functions

Overall, you may think of composition as a process of creating a *new* function from two existing ones.

However, our main concern is the challenge of identifying when a given function is a composite function.

For instance, given the function $y = (x + 1)^2$ to begin with, how can you *show* that it is a composite function?

One way is by explicitly defining the function's two component functions f and g , followed by demonstrating that the function can be written in the required form $g(f(x))$.

Identifying Composite Functions

Overall, you may think of composition as a process of creating a *new* function from two existing ones.

However, our main concern is the challenge of identifying when a given function is a composite function.

For instance, given the function $y = (x + 1)^2$ to begin with, how can you *show* that it is a composite function?

One way is by explicitly defining the function's two component functions f and g , followed by demonstrating that the function can be written in the required form $g(f(x))$.

Identifying Composite Functions

Overall, you may think of composition as a process of creating a *new* function from two existing ones.

However, our main concern is the challenge of identifying when a given function is a composite function.

For instance, given the function $y = (x + 1)^2$ to begin with, how can you *show* that it is a composite function?

One way is by explicitly defining the function's two component functions f and g , followed by demonstrating that the function can be written in the required form $g(f(x))$.

Identifying Composite Functions

Overall, you may think of composition as a process of creating a *new* function from two existing ones.

However, our main concern is the challenge of identifying when a given function is a composite function.

For instance, given the function $y = (x + 1)^2$ to begin with, how can you *show* that it is a composite function?

One way is by explicitly defining the function's two component functions f and g , followed by demonstrating that the function can be written in the required form $g(f(x))$.

Identifying Composite Functions (cont.)

So by setting $f(x) = x + 1$ and $g(u) = u^2$, we have:

$$\begin{aligned}y &= (x + 1)^2 \\&= (f(x))^2 && \longleftarrow \text{since } f(x) = x + 1 \\&= g(f(x)). && \longleftarrow \text{since } g(u) = u^2\end{aligned}$$

Since $f(x) = x + 1$ is 'nested' inside of $y = (x + 1)^2$, $f(x)$ is called the **inner** or **inside function**, and g is called the **outer** or **outside function**.

Identifying Composite Functions (cont.)

So by setting $f(x) = x + 1$ and $g(u) = u^2$, we have:

$$\begin{aligned}y &= (x + 1)^2 \\&= (f(x))^2 && \longleftarrow \text{since } f(x) = x + 1 \\&= g(f(x)). && \longleftarrow \text{since } g(u) = u^2\end{aligned}$$

Since $f(x) = x + 1$ is 'nested' inside of $y = (x + 1)^2$, $f(x)$ is called the **inner** or **inside function**, and g is called the **outer** or **outside function**.

Identifying Composite Functions (cont.)

So by setting $f(x) = x + 1$ and $g(u) = u^2$, we have:

$$\begin{aligned}y &= (x + 1)^2 \\&= (f(x))^2 && \longleftarrow \text{since } f(x) = x + 1 \\&= g(f(x)). && \longleftarrow \text{since } g(u) = u^2\end{aligned}$$

Since $f(x) = x + 1$ is 'nested' inside of $y = (x + 1)^2$, $f(x)$ is called the **inner** or **inside function**, and g is called the **outer** or **outside function**.

Identifying Composite Functions (cont.)

So by setting $f(x) = x + 1$ and $g(u) = u^2$, we have:

$$\begin{aligned}y &= (x + 1)^2 \\&= (f(x))^2 && \longleftarrow \text{since } f(x) = x + 1 \\&= g(f(x)). && \longleftarrow \text{since } g(u) = u^2\end{aligned}$$

Since $f(x) = x + 1$ is 'nested' inside of $y = (x + 1)^2$, $f(x)$ is called the **inner** or **inside function**, and g is called the **outer** or **outside function**.

Example

Here's another example. Consider the function $y = e^{3x}$.

Since the $3x$ appears to be 'nested' within the exponential function, we should set $f(x) = 3x$ and $g(u) = e^u$. Then:

$$y = e^{3x}$$

$$= e^{f(x)} \quad \longleftarrow \quad \text{since } f(x) = 3x$$

$$= g(f(x)). \quad \longleftarrow \quad \text{since } g(u) = e^u$$

Example

Here's another example. Consider the function $y = e^{3x}$.

Since the $3x$ appears to be 'nested' within the exponential function, we should set $f(x) = 3x$ and $g(u) = e^u$. Then:

$$y = e^{3x}$$

$$= e^{f(x)} \quad \longleftarrow \quad \text{since } f(x) = 3x$$

$$= g(f(x)). \quad \longleftarrow \quad \text{since } g(u) = e^u$$

Example

Here's another example. Consider the function $y = e^{3x}$.

Since the $3x$ appears to be 'nested' within the exponential function, we should set $f(x) = 3x$ and $g(u) = e^u$. Then:

$$\begin{aligned} y &= e^{3x} \\ &= e^{f(x)} && \longleftarrow \text{since } f(x) = 3x \\ &= g(f(x)). && \longleftarrow \text{since } g(u) = e^u \end{aligned}$$

Example

Here's another example. Consider the function $y = e^{3x}$.

Since the $3x$ appears to be 'nested' within the exponential function, we should set $f(x) = 3x$ and $g(u) = e^u$. Then:

$$\begin{aligned} y &= e^{3x} \\ &= e^{f(x)} && \longleftarrow \text{since } f(x) = 3x \\ &= g(f(x)). && \longleftarrow \text{since } g(u) = e^u \end{aligned}$$

Some More Examples (cont.)

Example

One more example. Consider the function $y = \sqrt{x^2 - 2}$.

Since $x^2 - 2$ appears to be 'nested' within the square root function, we should set $f(x) = x^2 - 2$ and $g(u) = \sqrt{u}$. Then:

$$\begin{aligned} y &= \sqrt{x^2 - 2} \\ &= \sqrt{f(x)} && \longleftarrow \text{since } f(x) = x^2 - 2 \\ &= g(f(x)). && \longleftarrow \text{since } g(u) = \sqrt{u} \end{aligned}$$

Example

One more example. Consider the function $y = \sqrt{x^2 - 2}$.

Since $x^2 - 2$ appears to be 'nested' within the square root function, we should set $f(x) = x^2 - 2$ and $g(u) = \sqrt{u}$. Then:

$$y = \sqrt{x^2 - 2}$$

$$= \sqrt{f(x)} \quad \longleftarrow \quad \text{since } f(x) = x^2 - 2$$

$$= g(f(x)). \quad \longleftarrow \quad \text{since } g(u) = \sqrt{u}$$

Some More Examples (cont.)

Example

One more example. Consider the function $y = \sqrt{x^2 - 2}$.

Since $x^2 - 2$ appears to be ‘nested’ within the square root function, we should set $f(x) = x^2 - 2$ and $g(u) = \sqrt{u}$. Then:

$$\begin{aligned} y &= \sqrt{x^2 - 2} \\ &= \sqrt{f(x)} && \longleftarrow \text{since } f(x) = x^2 - 2 \\ &= g(f(x)). && \longleftarrow \text{since } g(u) = \sqrt{u} \end{aligned}$$

Example

One more example. Consider the function $y = \sqrt{x^2 - 2}$.

Since $x^2 - 2$ appears to be ‘nested’ within the square root function, we should set $f(x) = x^2 - 2$ and $g(u) = \sqrt{u}$. Then:

$$\begin{aligned} y &= \sqrt{x^2 - 2} \\ &= \sqrt{f(x)} && \longleftarrow \text{since } f(x) = x^2 - 2 \\ &= g(f(x)). && \longleftarrow \text{since } g(u) = \sqrt{u} \end{aligned}$$

The process of verifying a given function as a composite function may seem laborious, but there is a good reason for developing this skill.

Suppose you wanted to differentiate

$$y = (x + 1)^2.$$

Expanding the right-hand side gives:

$$y = x^2 + 2x + 1,$$

and so

$$\frac{dy}{dx} = 2x + 2.$$

The process of verifying a given function as a composite function may seem laborious, but there is a good reason for developing this skill.

Suppose you wanted to differentiate

$$y = (x + 1)^2.$$

Expanding the right-hand side gives:

$$y = x^2 + 2x + 1,$$

and so

$$\frac{dy}{dx} = 2x + 2.$$

The process of verifying a given function as a composite function may seem laborious, but there is a good reason for developing this skill.

Suppose you wanted to differentiate

$$y = (x + 1)^2.$$

Expanding the right-hand side gives:

$$y = x^2 + 2x + 1,$$

and so

$$\frac{dy}{dx} = 2x + 2.$$

The process of verifying a given function as a composite function may seem laborious, but there is a good reason for developing this skill.

Suppose you wanted to differentiate

$$y = (x + 1)^2.$$

Expanding the right-hand side gives:

$$y = x^2 + 2x + 1,$$

and so

$$\frac{dy}{dx} = 2x + 2.$$

But how about incrementing the power from 2 to 7 and trying to differentiate

$$y = (x + 1)^7?$$

Unlike $y = (x + 1)^2$, one would need to expend considerably more effort in expanding the right-hand side:

$$y = x^7 + 7x^6 + 21x^5 + 35x^4 + 35x^3 + 21x^2 + 7x + 1.$$

Hence:

$$\frac{dy}{dx} = 7x^6 + 42x^5 + 105x^4 + 140x^3 + 105x^2 + 42x + 7.$$

But how about incrementing the power from 2 to 7 and trying to differentiate

$$y = (x + 1)^7?$$

Unlike $y = (x + 1)^2$, one would need to expend considerably more effort in expanding the right-hand side:

$$y = x^7 + 7x^6 + 21x^5 + 35x^4 + 35x^3 + 21x^2 + 7x + 1.$$

Hence:

$$\frac{dy}{dx} = 7x^6 + 42x^5 + 105x^4 + 140x^3 + 105x^2 + 42x + 7.$$

However, it turns out that there is a special technique called the **Chain Rule** (to be studied in Day 9) which can be used to differentiate a composite function *provided* its inside and outside functions have been identified explicitly.

For $y = (x + 1)^7$, this means recognising that if we set

$$u = x + 1,$$

we can then write

$$y = u^7.$$

However, it turns out that there is a special technique called the **Chain Rule** (to be studied in Day 9) which can be used to differentiate a composite function *provided* its inside and outside functions have been identified explicitly.

For $y = (x + 1)^7$, this means recognising that if we set

$$u = x + 1,$$

we can then write

$$y = u^7.$$

Practice Questions

Here are several practice questions to train your proficiency in decomposing composite functions.

For the following, identify the inside function u and express y as a function of u :

- ▶ $y = (x^3 + 5)^6$
- ▶ $y = e^{3-x^2}$
- ▶ $y = \sqrt{x^2 + 4x - 1}.$

Solutions

- ▶ $u = x^3 + 5, y = u^6;$
- ▶ $u = 3 - x^2, y = e^u;$
- ▶ $u = x^2 + 4x - 1, y = \sqrt{u}.$

- ▶ Given two functions f and g , the *composite function* “ g composed with f ” is defined to be the function $g(f(x))$.
- ▶ Decomposing a given composite function into its inside and outside functions is a prerequisite for using the *Chain Rule* for differentiation (studied the next day).